



Knowledge

***Clostridium perfringens* Type A: disease fundamentals and prevention strategies**

SUMMARY

Clostridium perfringens Type A is an emerging economic concern for dairy and beef producers because of its association with serious and often deadly gastrointestinal diseases in both cows and calves. This bacterium is a very versatile pathogen. It's ubiquitous in nature. Some forms of *C. perfringens* are found in all healthy livestock.

Many disease problems are caused by *C. perfringens* strains. The major toxins include alpha, beta, epsilon and iota, and all toxigenic types make alpha toxin. However, *C. perfringens*

Type A is the only one that makes **only** alpha toxin.

Clostridium perfringens Type A is associated with hemorrhagic bowel syndrome (HBS) in mature cows, which strikes apparently healthy animals without much warning and has a case fatality rate of 85 percent or higher. *Clostridium perfringens* Type A also is associated with abomasal ulcers, abomasal hemorrhage and abdominal tympany in calves, where it is frequently fatal.

Because treatment measures often are unsuccessful, prevention and management strategies are the best line of attack

against the disease. **Clostridium Perfringens Type A Toxoid** from Novartis Animal Health US, Inc. is the first and only vaccine for this disease to receive a conditional license by the USDA. The vaccine can be given to pregnant or non-pregnant animals to aid in the control of disease syndromes caused by the alpha toxin of *C. perfringens*. A Colorado State University study found that **Clostridium Perfringens Type A Toxoid** administered during the dry period led to increased anti-alpha toxin antibody in cows, as well as increased levels in their colostrum-fed calves.⁹

Clostridial organisms are commonly found in the environment and are part of the normal intestinal microflora of cattle.¹ However, dietary changes or parasitism may produce a favorable growth environment for these organisms, resulting in production of potent toxins that are harmful to the animal.

There are five types of *Clostridium perfringens* (A, B, C, D and E), which are identified by the main types of toxins they produce (alpha, beta, iota, epsilon and theta). *C. perfringens* Type A is identified as producing only alpha toxin out of the group of major toxins. Other toxins (beta 2 and enterotoxin) can be found in some isolates of *C. perfringens* Type A, but their role in disease is not known at this time. *C. perfringens* Type A and its alpha toxin are associated with serious gastrointestinal diseases in both young and mature animals.

Because treatment success is rare, emphasis is placed on preventative measures, such as vaccinating the pregnant cow to provide her with active immunity and provide her offspring passive immunity via colostrum.²

Until recently, there was not a commercial vaccine available for *C. perfringens* Type A, so producers relied on custom, or autogenous, vaccines to battle this clostridial strain. But now Novartis has **Clostridium Perfringens Type A Toxoid**, a conditionally licensed vaccine that gives veterinarians and producers a new tool in the battle against diseases associated with *C. perfringens* Type A.



C. perfringens Type A and mature cows

Hemorrhagic bowel syndrome (HBS), also known as jejunal hemorrhage syndrome or bloody gut, is a frustrating disease that strikes apparently healthy cattle without warning. HBS is more prevalent in dairy cows, especially in early lactation, although it also has been reported in beef cattle.³

Risk factors

In the last few years, several studies have indicated an association between HBS and *C. perfringens* Type A. While no single cause has been identified for HBS, *C. perfringens* is believed to be a contributor as it is commonly isolated from the gastrointestinal tracts of afflicted animals, while other enteric pathogens like *Salmonella* spp. and bovine viral diarrhea virus are rarely present in HBS cases.

Researchers have isolated *C. perfringens* Type A in both fecal cultures and blood clots in the jejunum of affected cows. A retrospective study from Colorado State University found that *C. perfringens* was isolated in 17 of 20 fecal samples and five of five intestinal biopsy specimens from cases of HBS.⁴

Signs of HBS

HBS begins with a sudden and sometimes massive hemorrhage into the small intestine, resulting in blood clots that obstruct the intestine. Although affected cows are often found dead or dying with no warning signs, following are some of the clinical symptoms of HBS:

- Sudden and complete anorexia
- Rapid pulse and respiratory rate
- Pale mucous membranes
- Severe decrease in milk production

- Severe depression
- Dark, tar-like feces, often containing clots of digested blood
- Abdominal distention, especially in lower right abdomen
- Normal or below-normal rectal temperature
- Scattered, low-pitched “pings” in lower right abdomen

The jejunum is the primary site of pathology, with findings including bloody fluid or clots in the lumen of the gut and a significant portion of the gut, up to three to four feet, affected.


Treatment of cattle with symptoms of HBS is rarely successful. Cattle may occasionally recover after treatment with fluids, laxatives, anti-inflammatory drugs and antibiotics, but more often, the disease progresses to ileus, intestinal necrosis, peritonitis and shock. Surgical treatment is often followed by complications of recurrent clotting and intestinal obstruction.

Prevention is best course

With a case fatality rate of 85 percent or higher, prevention must be the focus for HBS. In the past, prevention options have included the use of a custom or autogenous vaccine for *C. perfringens* Type A. Many veterinarians have reported favorable results from custom vaccines manufactured by Novartis.

In addition to vaccination, other management measures to consider are:

- Evaluating and correcting nutritional factors that may predispose cows to HBS, such as lack of roughage or too much carbohydrate in the diet
- Testing silage to verify proper ensiling, quality and to check for the presence of



Rod-shaped clostridial cells are apparent in the gram stain from the abomasal contents of a calf with distended abomasums due to clostridial abomasitis. Fluids from a healthy calf would normally have very few bacteria and no clostridia present.

C. perfringens Type A, mold or mycotoxins

- Practicing good bunk management, such as keeping feed pushed up and removing left-over feed from the bunk daily
- Identifying and correcting problems that might lower disease resistance during transition and early lactation



Ultrasound image of an HBS intestinal lesion.



Abdominal distention, especially in the lower right abdomen, is another symptom of HBS. When the animal is standing, the abdominal contour appears round or pear shaped.

To minimize the impact of *C. perfringens* Type A, producers should work with their veterinarian to determine the best vaccination and management protocols for their herds.

***C. perfringens* Type A and calves**

C. perfringens Type A also can be deadly for young beef and dairy calves. In calves, *C. perfringens* Type A is associated with abomasal ulcers, abomasal hemorrhage and abdominal tympany.

Common time periods to see *C. perfringens* in calves include:

- Less than one week of age
- Two months of age
- Seven to 20 weeks of age in veal calves

Calves show signs of quick onset of abdominal distension with pain, bloat, depression, feed refusal and sudden death. While there can be more than one cause for this syndrome, researchers have isolated *C. perfringens*

Type A from affected calves in several studies.²

For example, *C. perfringens* Type A was isolated from seven of eight neonatal calves that were referred to Kansas State University for clinical examination or necropsy. These animals – a mix of dairy and beef calves two to 21 days old – showed acute onset of abdominal tympany, colic, depression or death. Researchers concluded “the acute abdominal syndrome in the neonatal calves was unrelated to copper deficiency, and that *C. perfringens*, particularly Type A, may have had an appreciable contributory role in its pathogenesis.”⁵

In another study, intraruminal inoculation of *C. perfringens* Type A into healthy calves, four to 12 days old, induced anorexia, depression, bloat, diarrhea and in some cases, death.⁶

It is difficult to reproduce *C. perfringens* Type A disease in calves because it’s a multifactorial disease. Researchers haven’t found the complete complement of those factors in the case of calf disease. More research is needed to understand *C. perfringens* Type A.

If you suspect a case of *C. perfringens* Type A disease,

take the animal to a diagnostic facility. Diagnosis should:

- Document where the lesions are and the severity
- Include semiquantitative bacteriology culture for *C. perfringens* Type A
- Include genotyping of isolates
- Rule out infection by *C. perfringens* Types C and E

Standard clostridial seven-way vaccines do not protect against Type A, and until now, no commercial vaccines for *C. perfringens* Type A have been available. However, Novartis has produced custom, or autogenous, vaccines for *C. perfringens* Type A, and practitioners have been successful using them in herds where Type A has been isolated in calves.

***C. perfringens* Type A vaccination**

Novartis has been on the forefront of the *C. perfringens* Type A issue, conducting research and producing custom vaccines since the 1990s. After more than a decade of work, Novartis created the first USDA conditionally licensed vaccine – **Clostridium Perfringens Type A Toxoid for cattle.**



Prevention strategies are the best option for managing HBS. In addition to vaccination, there are nutritional management factors that can be employed.



***Clostridium perfringens* Type A can cause sudden death in young beef and dairy calves.**

The vaccine was tested using a study involving 13 vaccinates and four non-vaccinated controls. Vaccinates received two 2-mL SubQ doses of **Clostridium Perfringens Type A Toxoid** three weeks apart. Results showed that using the vaccine surpassed USDA requirements (i.e., 80 percent of animals demonstrating four international antitoxin units per mL), with eight international antitoxin units per mL being achieved in 89 percent of the vaccinated animals that were seronegative prior to vaccination. When including calves with

pre-vaccination titers, the seroconversion rate was 92 percent.⁷ (See Figure 1.)


Field studies involving 867 dairy and beef cattle – both open and pregnant – of various ages and breeds demonstrated the product is safe when used according to label directions.⁸

In addition, a Colorado State University study was conducted to determine if vaccinating pregnant cows and heifers with **Clostridium Perfringens Type A Toxoid** affected the serum neutralizing antibody titer to alpha toxin in the colostrum-fed calves of the vaccinated dams.⁹ Results found that the vaccine administered during the dry period led to increased anti-alpha toxin antibody in cows, as well as increased levels in their colostrum-fed calves.

Clostridium Perfringens Type A Toxoid is labeled for use in healthy cattle as an aid in the control of disease syndromes caused by the alpha toxin of *C. perfringens*. It contains a proprietary dual-component adjuvant system for optimum immune response and low tissue impact.

Two mL of the vaccine should be administered subcutaneously

in the neck, with revaccination in two to four weeks following the initial dose. Calves vaccinated prior to five months of age should be revaccinated at five to six months of age. Revaccinate annually or as recommended by your veterinarian. The product can be given to pregnant or non-pregnant animals, and has been proven safe in calves as young as one month of age.



1. Enterotoxemia Caused by *Clostridium perfringens* Type A. Merck Veterinary Manual. Eighth edition. 1998:445.
2. McGuirk S. Managing Clostridial Diseases in Cattle. *Proceedings of the 2002 Midwest Dairy Herd Health Conference*.
3. Abutarbush S, Carmalt J, Wilson D, O'Connor B, Clark E, Naylor J. Jejunal hemorrhage syndrome in 2 Canadian beef cows. *Can Vet J*. 2004;45:48-50.
4. Dennison A, VanMetre D, Callan R, Dinsmore P, Mason G, Ellis R. Hemorrhagic bowel syndrome in dairy cattle: 22 cases (1997-2000). *JAVMA*. 2002;221:686-689.
5. Roeder BL, Chengappa MM, Nagaraja TG, Avery TB, Kennedy GA. Isolation of *Clostridium perfringens* from neonatal calves with ruminal and abomasal tympany, abomasitis, and abomasal ulceration. *JAVMA*. 1987;190:1550-1555.
6. Roeder BL, Chengappa MM, Nagaraja TG, Avery TB, Kennedy GA. Experimental induction of abdominal tympany, abomasitis, and abomasal ulceration by intraruminal inoculation of *Clostridium perfringens* type A in neonatal calves. *Am J Vet Res*. 1988;49:201-207.
7. **Clostridium perfringens Type A Toxoid**. Code 8081.00. Report 1. Cattle Immunogenicity. Novartis Animal Health US Inc. February 3, 2004.
8. **Clostridium perfringens Type A Toxoid**. Code 8081.00. Report 5. Field Safety Testing. Novartis Animal Health US Inc. November 22, 2004.
9. Van Metre D. Passive transfer of neutralizing antibodies to *Clostridium perfringens* Type A alpha toxin in dairy calves fed colostrum from immunized dams. Paper presented at Annual Meeting of American Association of Bovine Practitioners 2006; Saint Paul, Minnesota.

Photos courtesy of Drs. Robert Callan and David Van Metre, College of Veterinary Medicine and Biomedical Sciences, Colorado State University.



www.livestock.novartis.com