

# PanSTAR™ Ery

Erysipelothrix Rhusiopathiae Bacterin

For use in healthy swine as an aid in the prevention of erysipelas caused by *Erysipelothrix rhusiopathiae* for a period of 24 weeks following vaccination.


## Product Numbers

### PanSTAR™ Ery

- 351 - 100 mL - 100 doses
- 352 - 250 mL - 250 doses

- Longest duration of immunity —** With 24 weeks duration of immunity (DOI), PanSTAR Ery has the longest, one-dose DOI claim with annual revaccination. See Chart 1.
- Low dose, easy to use —** 1-mL dose pig formulation; vaccinate pigs three weeks of age or older. Annual vaccination in breeding swine.
- Convenient —** Available in 100-, 250- and 500-dose bottles. See Table 1.

**Chart 1. Percent Protection 24 Weeks After Vaccination**



Product photo

**Table 1. Number Swine Protected at Various Times Post-vaccination**

	5 weeks	9 weeks	13 weeks	17 weeks
<b>One Dose</b>	4 of 4	5 of 5	4 of 5	5 of 5
<b>Two Dose*</b>	4 of 4	5 of 5	4 of 4	4 of 5
<b>Controls</b>	0 of 4	0 of 4	0 of 4	0 of 5

One-month-old pigs were vaccinated with either one dose or two doses of PanSTAR™ Ery and experimentally challenged at various times post-vaccination. A single dose gave the same protection as the two-dose program.

\*Second dose not required in the field. Study was done for experimental purposes.


## PanSTAR™ Ery

**ADJUVANT:** ImmunSTAR™

**DIRECTIONS:** Shake well before using. Administer 1 mL intramuscularly at three weeks of age or older.

**PRECAUTIONS:** Store out of direct sunlight at 2°-7° C (35°-45° F). DO NOT FREEZE. Use entire contents when first opened. Do not vaccinate within 60 days prior to slaughter. Anaphylactic reactions may occur. Symptomatic treatment: Epinephrine.

**PACKAGING:** 100 mL/100 dose and 250 mL/250 dose



**Customer Service**  
**(800) 843-3386**

[www.livestock.novartis.com](http://www.livestock.novartis.com)  
© 2003 Novartis Animal Vaccines, Inc.  
Printed in USA  
4504998, OCT03